

SHARKOV, V. I.

Contribution of science to the technological progress of the
hydrolysis industry. Gidroliz. i lesokhim. prom. 14 no.6:10-13
'51. (MIRA 14:9)
(Hydrolysis)

SHARKOV, V.I.; LEVANOVA, V.P.; ARTEM'YEVA, I.S.

Packing density of some natural holocellulose, Zhur.prikl.khim.
34 no.11:2508-2515 N '61. (MIRA 15:1)

1. Gosudarstvennyy nauchno-issledovatel'skiy institut gidroliznoy
i sul'fitno-spirtovoy promyshlennosti.
(Holocellulose)

SHARKOV, V.I.; SAPOTNITSKIY, S.A.

Solve the problem in the best interests of the state. Bum.prom. 36
no.2:9 F '61. (MIRA 14:2)

1. Direktor Nauchno-issledovatel'skogo instituta gidroliznoy i
sul'fitno-spirtovoy promyshlennosti (for Sharkov). 2. Zaveduyushchiy
laboratoriyye pereabotki sul'fitnykh shchelokov Nauchno-issledovateli'-
skogo instituta gidroliznoy i sul'fitno-spirtovoy promyshlennosti
(for Sapotnitskiy).

(Woodpulp)

(Alcohol)

SHARKOV, V.I.; LEVANOVA, V.P.; BOLOTOVA, A.K.

Supermolecular structure of extrastrong cellulose hydrate
fibers. Khim.volok. no.5:32-36 '62. (MIRA 15:11)
(Textile fibers, Synthetic)
(Cellulose)

SHAROV, V. I., KUBINA, N. S., SLOV'YANA, Yu. N.

Properties and isolation of glucosaminan from wood
hemicellulose of fir (*Picea excelsa*). Zhur. prikl. khim.
V. no. 11119. 1128 by 62. (MFA 15:5)
(Glucosaminane)
(Fir)

TOKAREV, B.I.; SHARKOV, V.I.

Determining oxymethylfurfurole in wood hydrolyzates. Gidroliz.i
lesokhim.prom. 15 no.6:16 '62. (MIRA 15:9)
(Wood--Chemistry) (Furaldehyde)

SLAVYANSKIY, Aleksey Konstantinovich, prof.; SHARKOV, Vasiliy Ivanovich, prof.; LIVEROVSKIY, Aleksey Alekseyevich, dots.; BUYEVSKOY, Anatoliy Vasil'yevich, dots.; MEDNIKOV, Fedor Alekseyevich, dots.; LYAMIN, Vladimir Aleksandrovich, dots.; SOLODKIY, Fedor Timofeyevich, dots.; TSATSKA, Elio Mat'-Iudovich, dots.; DMITRIYEVA, Ol'ga Andreyevna, assistant; NIKANDOROV, Boris Fedorovich, inzh.; GORDON, L. V., kand. tekhn. nauk, retsenzent; SUKHANOVSKIY, S. I., red.; KHOT'KOVA, Ye. S., red.izd-va; SHIBKOVA, R. Ye., tekhn. red.

[Chemical technology of wood] Khimicheskaiia tekhnologija drevesiny. Moskva, Goslesbumizdat, 1962. 574 p. (MIRA 16:4)
(Wood—Chemistry)

YEFIMOV, V.A.; MOLCHANOV, M.N.; SHARKOV, V.I.

Treatment of sunflower seed shells in a continuous horizontal pilot hydrolyzer. Sbor.trud. NIIGS 11:16-22 '63. (MIRA 16:12)

KRUPNOVA, A.V.; SHARKOV, V.I.

Thermomechanical method for converting cellulose into a readily hydrolyzable state. Gidroliz. i lesokhim.prom. 16 no.3:8-10 '63. (MIRA 16:5)

1. Gosudarstvennyy nauchno-issledovatel'skiy institut gidroliznoy i sul'fitnospirtovoy promyshlennosti.
(Cellulose) (Hydrolysis)

SHARKOV, V.I.; SHCHEGOLOVA, T.A.

Obtaining cellulose preparations with different density of packing
of the macromolecules. Zhur.prikl.khim. 36 no.6:1326-1330 Je
'63. (MIRA 16:8)
(Cellulose)

TOKAREV, B.I.; SHARKOV, V.I.

Behavior of hydroxymethylfurfurol during alcoholic fermentation
of hydrolyzates. Gidroliz. i lesokhim. prom. 16 no.2:4-5 '63.
(MIRA 16:6)
(Furaldehyde) (Glucose) (Fermentation)

SAPOTNITSKIY, S.A.; MYASNIKOVA, R.M.; SHARKOV, V.I.

Use of SO₂ for the inversion of oligosaccharides in the liquor
of bisulfite cooking of vegetable raw materials. Gidroliz. i
lesokhim.prom. 17 no.1:12-13 '64. (MIRA 17:4)

1. Gosudarstvennyy nauchno-issledovatel'skiy institut gidroliznoy
i sul'fitno-spirtovoy promyshlennosti.

US-ANOV, Kh.U.; NIKONOV, V.S.; ZARIPOVA, A.M.; SHABLOV, V.I.,
doktor tekhn.nauk, prof., otv. red.; SOKOLOVA, A.A., red.

[Prospects of the chemical processing of cotton waste] Per-
spektivy khimicheskoi pererabotki otkhodov khlopkovodstva.
Tashkent, Izd-vo "Nauka" UzSSR, 1964. 125 p.
(MIRA 17:11)

KRUPNOVA, A.V.; SHAROV, V.I.

Mechanical and thermochemical method of transforming cellulose-containing materials to a readily hydrolyzable state. Gidrolyz. i lesokhim. prom. 17 no.3:3-5 '64. (MIRA 17:9)

1. Gosudarstvennyy nauchno-issledovatel'skiy institut gidrolyznoy i sul'fitno-spirtovoy promyshlennosti.

DMITRIYEVA, O.A.; POTAPOVA, N.P.; SHARKOV, V.I.

Comparative study of the supermolecular structure of wood
cellulose by the methods of thermal and hydrolytic action.
Zhur. prikl. khim. 37 no. 6:2083-2085 S '64.

(MIRA 17:10)

VYRODOVA, L.P.; SHAROV, V.I.

Studying the dissolving process of cellulose and wood in concentrated sulfuric acid. Sbor. trud. NIIGS 12:40-48 '64.

(MIRA 18:3)

BOLOTCEVA, A.K.; SHARKOV, V.I.

Using the method of dielectric constant measurement in the study
of the supramolecular structure of cellulose. Sbor. trud. NII GS
12:49-59 '64.

Investigating the capillary structure of cellulose. Ibid.:60-70

Retarding effect of water in the hydrolysis reaction of cellulose.
Ibid.:71-86 (MIRA 18:3)

MITAIVINA, O.A.; POTAPOVA, N.P.; SHARKOV, V.I.

Comparative study of cotton cellulose supermolecular structure
by the methods of thermal and hydrolytic treatment. Tur. prikl.
Khim. 37 no.7:1583-1589 J1 '64. (MIRA 18:4)

1. Leningradskaya lesotekhnicheskaya akademiya v. v. Kirova i
Gosudarstvennyy institut gidrolyznoy promyslennosti.

SHAPROV, V. V., VILNIUS, M.L., MO-GU/HEVIA, YU.F.

Study of the homogeneity of xyluronide by fractionation. (Bur. prikl. khim. 38 no. 701525-13/3 31 165. (MIR: 1817)

1. Gaudans'vianyy nauchno-issledovatel'skiy institut giproliancy i prikladnoi tekhniki po promyshlennosti.

SHARKOV, V. M.

USER/ Engineering - Industrial processes

Card 1/1 Pub. 103 - 5/19

Authors : Sokolovskiy, M. V.; Gumenyy, V. N.; and Sharkov, V. M.

Title : Thermal treatment of worm threads with high-frequency currents

Periodical : Stan. i instr. 2, 19 - 20, Feb 1955

Abstract : The construction of a special machine for hardening of worm threads with high-frequency current was announced by the "Krasniy Metallist Metallurgical Plant." The technical and mechanical properties of the new machine are described. It was found that the changes in the worm dimension after thermal treatment with high-frequency currents are very insignificant and can be totally disregarded. The thermal treatment cycle of the new machine is 15 - 18 times smaller than otherwise and the mechanical properties of the treated part remain unchanged. Drawings.

Institution:

Submitted:

SHARKOV, V. M.

"The effective technique of prospecting useful minerals in less investigated regions"

report to be submitted to the United Nations Conference on the
Application of Science and Technology for the Benefit of the Least
Developed Countries, Addis Ababa, 4-20 Feb 63

SHARKOV, V.N.

The AK-1T industrial power truck. Biul.tekh.-ekon. inform.
no.3:65-66 '61. (MIRA 14:3)
(Industrial power trucks)

U.S.S.R., 1950.

"Experience of Utilizing aeromethods in Geological-geomorphological Research in
Conditions of the Golodnaya Steppe of Kazakhstan," Trudy Lab. Aeromethods,
Vol. II, p. 93, 1950.

YAKOVLEV, S.A.; APUKHTIN, N.I.; BOCH, S.G.; VOZNESENSKIY, D.V.; GROMOV, V.I.; ZHUKOV, M.M.; KRASNOV, I.I.; LUNGERGAUZEN, G.F.; PERKONS, V.A.; POKROVSKAYA, I.M.; RUDOVITS, Yu.L. [deceased]; STAGNOVA, A.S.; SHARKOV, V.V.; EPSHTEYN, S.V.; YAKOVLEVA, S.V.; VERSTAK, G. V. ~~redaktor~~; GUDOV, O.A., tekhnicheskiy redaktor.

[Methodical aid for studying and geological surveying of quaternary deposits; description of methods] Metodicheskoe rukovodstvo po izucheniiu i geologicheskoi s"emke chetvertichnykh otlozhenii; opisanie metodov. Sost. S.A. Iakovlev. Moskva, Gos. nauchno-tekhn. izd-vo lit-ry po geologii i okhrane nedr, 1955. 485 p. [Microfilm] (MLRA 9:1)

1. Leningrad. Vsesoyuznyy geologicheskii institut.
(Geological surveys) (Geology, Stratigraphic--Quaternary--
Study and teaching)

Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 12, 15-1957-12-16943
p 40 (USSR)

AUTHOR: Sharkov, V. V.

TITLE: Use of Aerial Methods in the Studies of Quaternary Deposits (Primeneniye aerometodov pri issledovaniyakh chetvertichnykh otlozheniy)

PERIODICAL: V sb: Metod. rukovodstvo po izucheniyu i geol. s "yemke chetvertich. otlozheniy, ph 2, Moscow, Gosgeoltekhnizdat, 1955, pp 56-78

ABSTRACT: Bibliographical entry

Card 1/1

SHAKHLEV, V. V.

Example of using aerial photographs to show changes in the actual thickness of ocean rock bottom. Trudy Lab. aeromet. 4:138-140 '55.
(Submarine geology) (Photography, Aerial) (MLRA 9:2)

GUR'YEVA, Z.I.; SHARKOV, V.V.; SHUL'TS, S.S.

Results of geological mapping of shallow ocean bottom areas by
means of aerial photographs. Sov.geol. no.42:65-79 '55.

(MLRA 8:6)

(Photography, Aerial) (Geology--Maps)

SHARKOV, Vitaliy Vasil'evich, redaktor; KUDRITSKIY, Dmitriy
Mikhaylovich, redaktor; DOLMATOV, P.S., vedushchiy redaktor;
GRINAD'YEVA, I.M., tekhnicheskiy redaktor

[Use of aerial methods for geological investigations of the
ocean bottom; a brief practical manual] Primenenie aerometodov
dlia geologicheskikh issledovanii morskogo dna; kratkoe
metodicheskoe rukovodstvo. Pod red. V.V. Sharkova i D.M.
Kudritskogo. Leningrad, Gos. nauchno-tekhn. izd-vo neftianoi i
gorno-toplivnoi lit-ry, Leningradskoe otd-nie, 1956.
254 p.

(MLRA 10:4)

1. Akademiya nauk SSSR. Laboratoriya aerometodov.
(Aerial photogrammetry) (Ocean bottom)

SHARKOV, V.V., kand.geogr.nauk, otvetstvennyy red.; ARON, G.M., red.izd-va;
BLEYKH, B.Yu., tekhn.red.

[Geological air survey of shallow parts of the Caspian Sea]
Aerogeologicheskaiia s"emka melkovodnykh zon Kaspiiskogo moria.
Moskva, 1958. 139 p.
(MIRA 11:6)

1. Akademiya nauk SSSR. Laboratoriya aerometodov.
(Caspian Sea--Aerial photogrammetry)

3(4) PHASE I BOOK EXPLOITATION SOV/1835

Akademiya nauk SSSR. Laboratoriya aerometodov

Trudy, t. 6 (Transactions of the Laboratory of Aerial Methods, USSR Academy of Sciences, Vol 6) Moscow, Izd-vo AN SSSR, 1958. 280 p. Errata slip inserted. 1,500 copies printed.

Resp. Ed.: V.P. Miroshnichenko, Candidate of Geological and Mineralogical Sciences; Ed. of publishing House: D.M. Kudritskiy; Tech. Ed.: E.Yu. Bleykh.

PURPOSE: This volume is intended for geologists, photo interpreters, or other personnel engaged in the study of landscape formations, especially from the standpoint of aerial photography.

COVERAGE: This collection of studies and brief articles treats problems in aerial photography and photo interpretation in relation to geological phenomena. The geographical area of study, with minor exceptions, is the Caspian plains and western shore. Most of the studies are well illustrated with aerial photographs. Aside from the numerous articles on geological phenomena of the Caspian basin, the following are also covered: portions of the Russian platform, the Muyunkumy sands of Central Kazakhstan, photo interpretation of clayey flats, desert vegetation and tree cover, the effective lens speed of photographic objectives, photogrammetric determination of profiles on hydro technical models, and others. No personalities are mentioned. References follow each main article.

TABLE OF CONTENTS:

Our'yeva, Z.I. The Origin of Some "Cove-forming" Types of Relief on the Western Littoral of the Caspian Sea	234
Sharkov, V.V. The Violent Eruption of Gas on the Caspian Sea Bottom, Northeast of Cape Aniye	241
Our'yeva, Z.I., and V.V. Sharkov. Some Characteristics of the Geomorphological Structure of the Western Coast of the Caspian Sea from Cape Kilyazinskaya Bar to the Mouth of the Samur River	243
Kharin, N.O. Experience in the Use of Aero Visual Observations in Studying Forests	

GUR'YEVA, Z. I.; SHARKOV, V.V.

Geomorphological characteristics of the structure of the Caspian
western shore from the Kilyazi spit to the Samur estuary. Trudy Lab.
aeromet. 6:243-256 '58. (MIKA 12:1)
(Caspian Sea--Shore lines)

SHARKOV, V.V.

Use of aerial methods in solving some problems in connection with
the lowering level of the Caspian Sea. Trudy Okean, kom. 5:202-
207 '59. (MIRA 13:6)
(Caspian Sea--Hydrographic surveying) (Aerial photogrammetry)

PHASE I BOOK EXPLOITATION

SOV/4315

SOV/7-S-9

Akademiya nauk SSSR. Laboratoriya aerometodov

Trudy, tom 9 (Transactions of the Laboratory of Aerial Methods, USSR Academy of Sciences, vol. 9) Moscow, AN SSSR, 1960. 357 p. Errata slip inserted. 1,700 copies printed.

Resp. Ed.: V.V. Sharkov, Candidate of Geography; Ed. of Publishing House: D.M. Kudritskiy; Tech. Ed.: M.Ye. Zendel'.

PURPOSE: This volume is intended for geographers, geologists, geodesists, and photogrammetrists.

COVERAGE: This collection of 23 articles contains studies of the earth's surface, structure, and geological formations by means of aerial photography. The authors discuss the principles, methods and techniques used in aerial surveying to determine such factors as the petrographic composition of the soil through the measurement of the spectral brightness of surfaces, the geological structure of underwater areas through recorded photographic images, the geological composition and geomorphological structure of underlying layers through the analysis of surface plant coverings, the trends and characteristics of recent tectonic movements through the study of surface features traced photographically

Card 1/5

Transactions of the Laboratory (Cont.)

SOV/4315

over extensive regions, etc. The instruments used in this work (cameras, cinematographic spectrographs, stereographs, stereoprojectors, color and black and white film) are described and the analysis and interpretation of the data obtained discussed. References accompany individual articles.

TABLE OF CONTENTS:

Miroshnichenko, V.P. "Takyrs" as Indicators of the Most Recent Tectonic Movements in the Sandy Deserts of Central Asia	3
Romanova, M.A. Experimental Aerial Petrographic Survey of the Sands of the Northwestern Caspian Region	40
Gur'yeva, Z.I., and V.V. Sharkov. Geologic Structure of the Underwater Slope of the Southwestern Part of the Taman' Peninsula	82
Tolchel'nikov, Yu.S. Natural Factors Affecting the Tone of the Soil Images of Plowed Massifs on Aerial Photographs	101

Card 2/5

SHARKOV, V.V.

Aerial methods of studying outcrops of iron ore on the submarine
slope of the Black Sea south of the Taman Peninsula. Trudy Lab.
aeromet. 10:10-14 '60. (MIRA 14:1)
(Black Sea—Iron ores) (Aeronautics in geology)

SHARKOV, V.V.; GUR'YEVA, Z.I.

Some examples of the use of aerial photographs in detecting
anticlinal structures. Trudy Lab. aeromet. 10:15-23 '60.

(Aeronautics in geology) (Photography, Aerial) (MIRA 14:1)

SHARKOV, V.V.; GUR'YEVA, Z.I.; KOSHECHKIN, B.I.

Some features of the geological structure of the submarine slope
of the taman Peninsula in the Sea of Azov (according to the
materials of aerogeological research). Trudy Lab. aeromet.
10:24-34 '60. (MIRA 14:1)
(Azov, Sea of—Submarine geology)

SHARKOV, V. V., Doc GEOL ~~AND MINERAL~~ Sci, "GEOLOGICAL
STRUCTURE AND HISTORY OF THE DEVELOPMENT OF THE RELIEF
OF THE CONTINENTAL SLOPE OF THE WEST ^{AND} COAST OF THE CASPIAN
SEA, ^{AND} FROM THE CITY OF MAKHACHKALA TO THE ESTUARY OF THE
KURA RIVER (ON THE BASIS OF AN AERIAL PHOTOGRAPHIC SURVEY).
LENINGRAD, 1961. (ALL-UNION SCI RES GEOL INST "VSEGYEI").
(KL-DV, 11-61, 212).

-61-

SHARKOV, V.V.; GUR'YEVA, Z.I.

Geomorphology of the continental slope of the Taman' Peninsula.
Uch.zap.LGU no.298:155-170 '61. (MIRA 15:2)
(Taman' Peninsula--Slopes (Physical geography))

S/035/62/000/011/058/079
A001/A101

AUTHORS: Sharkov, V. V., Gur'yeva, Z. I.

TITLE: An experience in studying the landscapes of shallow sea floor sections in the western coast of the Caspian Sea and Caucasian-Taman' coast of the Black Sea

PERIODICAL: Referativnyy zhurnal, Astronomiya i Geodeziya, no. 11, 1962, 19, abstract 11G144 (In collection: "Primeneniye aerometodov v landshaftn. issled.", Moscow - Leningrad., AN SSSR, 1961, 255 - 277)

TEXT: Aerial photosurvey of sea floor was experimentally performed with AFA (AFA) middle-focus cameras on films of types, P Φ -1 (RF-1) and RF-3, panchromatic film of type 10-600, multi-layer colored one, etc. Best results were obtained in photographing with yellow light filter on the film RF-1 and RF-3. The following scales were used: 1 : 25,000; 1 : 10,000 and 1 : 5,000 (on small areas 1 : 3,000 and 1 : 2,000). It was found that in surveys on scales finer than 1 : 10,000 the sea floor is difficult to decipher, as many objects are not reproduced on photographs. Field deciphering was made by running routes on dry

Card 1/2

An experience in studying the...

S/035/62/000/011/058/079
A001/A101

land along the shore and series of tacks into the sea in a cutter and boat. The sea floor was studied by means of lifting samples with devices, and in main sections scientific workers carried out diving immersions. The floor relief was studied by means of a P0J -1M (REL-1M) sounding device. The authors consider briefly the signs for deciphering objects of sea floor, established as a result of experimental work, which enable one to identify on aerial photographs outcrops of various primary rocks to the sea floor and recent deposits, to determine their composition and relative geologic age, and sometimes to determine even the character of rock occurrence, individual elements of faults, breaks, and to detect the existence of different genetic relief forms, vegetation, etc. There are 16 references.

R. Vol'pe

[Abstracter's note: Complete translation]

Card 2/2

SHARKOV, Vitaliy Vasil'yevich; MOROZOV, A.I., otv. red.

[Geology of the underground slope of the western coast of the Caspian Sea (from the city of Makhachkala to the Kura estuary)] Geologija podvodnogo sklona zapadnogo beregа Kaspiiskogo moria (ot g. Makhachkala do ust'ia r. Kury). Moskva, Izd-vo "Nauka," 1964. 429 p.

(MIRA 17:6)

SFARKOV, Ye.

Members of the scientific technological society aid a collective farm. Mashinostroitel' no.11:47 N '61. (MIRA 14:11)
(Kalinin Province--Collective farms)

"APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R001548620004-1

Re: [REDACTED] G-1000, 1970

Re: [REDACTED] Standardized [REDACTED] 1970
(MIA: 1F:7)

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R001548620004-1"

SHAKOV, Ye.V.

Evolution of rock-forming minerals in the process of creating
the Zlatogorsk differentiated pluton (northern Kazakhstan).
Vest. LGU 20 no.24:37-41 '65. (MFA 19:1)

1. Submitted May 15, 1965.

SHARKOV, Yu.

Nature of land income in the agricultural cooperatives of
socialist countries. Vop. ekon. no.1:151-154 Ja '64.
(MIRA 17:3)

SHARKOV, Yu.V.; GLIKO, O.A.

Geochemical methods of prospecting for ore deposits in foreign countries. Razved. i okh.nedr 22 no.3:55-62 Mr '56.(MIRA 9:7)
(Geochemical prospecting)

SHARKOV, Yu.V.

Peculiarities of conducting trace-metal prospecting under varying natural conditions. Razved.i okh.nedr. 22 no.8:15-22 Ag '56.
(MLRA 9:11)

1. Tekhupravleniye Ministerstva geologii i okhrany nedr SSSR.
(Geochemical prospecting) (Ore deposits)

KRASHNIKOV, V.I., glavnnyy red.; BRODSKIY, A.A., red.; PEREL'MAN, A.I., red.;
SAUKOV, A.A., red.; SAFRONOV, N.I., red.; SERGEYEV, Ye.A., red.;
KHITAROV, N.I., red.; SHARKOV, Yu.V., red. SHCHERBINA, V.V., red.;
GUROVA, O.A., tekhn.red.

[Geokhimicheskie poiski rudnykh mestorozhdenii v SSSR; trudy soveshchaniia. Pod red. V.I.Krasnkova. Moskva, Gos.nauchno-tekhn.izd-vo lit-ry po geol. i okhrane nedr, 1957. 466 p. (MIRA 11:3)

1. Vsesoyuznoye soveshchaniye po geokhimicheskim metodam poiskov
rudnykh mestorozhdeniy. 1st, Moscow, 1956.
(Geochemical prospecting)

SOV/26-59-12-3/45

MATERIAL:

Sharkov, Yu. V., Candidate of Geological and Mineralogical Sciences (Moscow)

-- 100

JOURNAL:

Searching for Deposits of Atomic Raw Material

SUBJECT:

Priroda, 1959, Nr 12, pp 13-21 (USSR)

The author states that the splitting of the atom has created a great demand for uranium. Most uranium deposits presently being worked are those which were easiest to find. As they become exhausted, the problem is to find others. The article is an account of methods of prospecting employed in the USSR. To enable large areas to be surveyed easily and quickly, radiometers and counters have been developed. The instruments and methods used for prospecting from aircraft are described. The author says that aircraft have been used for this purpose in the USSR since 1947. Light MI-1 and KA-15 helicopters, carrying a 45-kg scintillometer, are used. Practice

Page 1/3

-- certain natural conditions

SOV/26-59-12-3/45

Searching for Deposits of Atomic Raw Material

is described. There are 3 photographs, 2 diagrams,
and 1 chart. *✓*

Card 3/3

KRASNIKOV, V.I.; SHARKOV, Yu.V.

Spatial and genetic relation between exogenetic and sedimentary-metamorphogenetic uranium deposits and arid zones of the corresponding geologic epochs. Dokl. AN SSSR. 144 no.6: 1359-1362 Je '62. (MIRA 15:6)

1. Predstavлено акад. D.I. Shcherbakovym.
(Uranium ores)

SHARKOV, Yu. V.

"Methodological instructions for carrying out geochemical prospecting for pyritic copper deposits in the Southern Urals based on the secondary halos of dispersion" by M. A. Glazovskia. Reviewed by Yu. V. Sharkov. Vop. geog. no. 59:178-179 '62.
(MIRA 16:1)

(Ural Mountains—Geochemical prospecting)
(Ural Mountains—Chalcopyrite)
(Glazovskia, M. A.)

SAUKOV, A.A.; GINZBURG, I.I.; PEREL'MAN, A.I.; AYDIN'YAN, N.Kh.;
SHARKOV, Yu.V.

Vladimir Ivanovich Krasnikov; obituary. Geol. rud. mestorozh.
5 no.2:141-142 Mr-Ap '63. (MIRA 16:6)

(Krasnikov, Vladimir Ivanovich, 1907-1962)

KRAVIKOV, Vladimir Ivanovich (1906-1962), prof., doktor geol.-miner. nauk; DYUKOV, A.I., otv. red.; KAZHDAN, A.B., otv. red.; PEREL'MAN, A.I., red.; SHERKOV, Yu.V., red.

[Fundamentals of an efficient method of prospecting for ore deposits] Osnovy ratsional'noi metodiki poiskov rudnykh mestorozhdenii. 2. izd. Moskva, Nedra, 1965. 398 p.
(MIRA 18:12)

SHARKOVA, A. S.

DECEASED

1964

MICROBIOLOGY

c/1963

SHARKOVA, K.D.

Result of joint work of parasitological departments and prophylactic disinfection departments in Stalingrad. Med.paraz. i paraz.bol. 27 no.6:728-729 N-D '58. (MIRA 12:2)
(PARASITIC DISEASES, prev. & control,
cooperation of parasitol. & disinfection
departments (Rus))
(ANTISEPSIS AND ASEPSIS,
same)

KUCHEROVA, N.F.; ZHUKOVA, I.G.; KAMZOLOVA, N.N.; PETRUCHENKO, M.I.;
SHARKOVA, N.M.; KOCHETKOV, N.K.

Indole derivatives. Part 8:9-Acyl-1,2,3,4, 4a, 9a-hexahydro-8-carbolines. Zhur. ob. khim. 31 no. 3:930-936 Mr '61, (MIRA 14:3)

1. Nauchno-issledovatel'skiy institut farmakologii i khimioterapii.
(Pyridindole)

SHARKOVA, N.M.; KUCHEROVA, N.F.; ZAGOREVSKIY, V.A.

Derivatives of indole. Part 9: Synthesis of derivatives
of pyrano (4,3-b)-3,4-dihydroindoles and chromeno (4,3-b)indoles.
Zhur.ob.khim. 32 no.11:3640-3645 N '62. (MIRA 15:11)

1. Institut farmakologii i khimioterapii AMN SSSR.
(Pyranoindole) (Benzopyranoindole)

SHARKOVA, N.M.; EUCHIROVA, N.F.; ZAGOREVSKIY, V.A.

Derivatives of indole. Part 15: Syntheses of some condensed indoline systems. Zhur. ob. khim. 34 no. 5:1614-1618
My '64. (MIRA 17:7)

1. Institut farmakologii i khimioterapii AMN SSSR.

AUTHOR: Sharkova, N.V.

SOV/140-58-3-33/34

TITLE: The Application of the Method of Oscillating Functions for the Approximative Solution of Initial and Boundary Value Problems for Non-Linear Differential Equations of Second Order (Primeneniye metoda otsilliruyushchikh funktsiy k priblizhennomu resheniyu nachal'nykh i krayevykh zadach dlya ne-lineynykh differentsiyal'nykh uravneniy vtorogo poryadka)

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy. Matematika, 1958, Nr 3, pp 251-255 (USSR)

ABSTRACT: The method proposed by Mel'nikov [Ref 1-3] for the approximative solution of differential equations of first order is extended to the solution of $y'' = f(x, y, y')$. The solution of the initial value problem $y(0) = a$, $y'(0) = b$ is sought in the form

$$y_N(x) = \frac{a_k}{2} (x - x_k)^2 + b_k (x - x_k) + c_k, \quad x_k \leq x \leq x_{k+1},$$

whereby the initial conditions, the continuity of the solution etc. are used for the determination of the coefficients. The substitution of the obtained approximation into (1) then leads to oscillating functions which can be estimated more or less

Card 1/2

The Application of the Method of Oscillating Functions for the Approximative Solution of Initial and Boundary Value Problems for Non-Linear Differential Equations of Second Order

SOV,140 -58-3-33/34

roughly.

The boundary value problem $y(0) = y(1) = 0$ is solved by enclosing the solution between the solutions of two initial value problems $[y(0) = 0, y'(0) = b_1]$ and $[y(0) = 0, y'(0) = b_2]$

whereby the b_1 and b_2 depend on the errors of the approximative solution of the initial value problems.

There are 4 references, 3 of which are Soviet, and 1 German.

ASSOCIATION: Permskiy gosudarstvennyy universitet imeni A.M.Gor'kogo
(Perm' State University imeni A.M.Gor'kiy)

SUBMITTED: December 20, 1957

Card 2/2

16,3400

32472
S/044/61/000/010/038/051
C111/C222AUTHOR: Sharkova, N.V.

TITLE: The application of the method of oscillating functions for the approximate solution of boundary value problems for differential equations of second order

PERIODICAL: Referativnyy zhurnal. Matematika, no. 10, 1961, 36-37.
abstract 10 V 222. ("Uch. zap. Permsk. un-t", 1959, 13.
no. 2, 41-61)

TEXT: The method of oscillating functions developed in the papers of S.I. Mel'nik (R Zh Mat, 1954. 5256 ; 1957, 5942, 5943) is used for the approximate solution of the linear differential equation of second order

$$y'' = p(x)y' + q(x)y + r(x)$$

with the boundary conditions

$$R_1[y] \equiv \alpha_1 y(0) + \alpha'_1 y'(0) + \beta_1 y(1) + \beta'_1 y'(1) = 0 ,$$

$$R_2[y] \equiv \alpha_2 y(0) + \alpha'_2 y'(0) + \beta_2 y(1) + \beta'_2 y'(1) = 0 ,$$

Card 1/2

The application of the method of ...

32472
S/044/61/000/010/038/05,
C111/C222

where $p(x)$, $q(x)$ and $r(x)$ are functions differentiable on $[0,1]$, and the rank of the matrix

$$\begin{pmatrix} \alpha_1 & \alpha'_1 & \beta_1 & \beta'_1 \\ \alpha_2 & \alpha'_2 & \beta_2 & \beta'_2 \end{pmatrix}$$

X

equals 2. An estimation of the approximate solution is given (under the assumption that the estimation for the Green's function is known). For the equation $y'' = q(x)y + r(x)$ with the conditions $y(0) = y(1) = 0$ the error estimation can be obtained without a Green's function. The method of oscillating functions is used for the determination of the eigenvalues of boundary value problems and for the approximate solution of the Cauchy and the boundary value problem (under some restrictions) for the non-linear differential equation $y'' = f(x, y, y')$. Examples are considered.

[Abstracter's note : Complete translation.]

Card 2/2

Shchukin, V. I., et al. of Phys-Math Sci -- (disc) "Use of Oscillatory Functions for the Solution of Regional Problems for Common Differential Equations," Perm', 1959, 15 pp (Perm' State Univ im A. M. Gor'kiy) (KL, 2-63, 110)

S/140/60/000/003/011/011
C111/C222

16,3400

AUTHOR: Sharkova, N.V.TITLE: Approximate Solution of Differential Equations With a Lagging Argument With the Aid of the Method of Oscillating FunctionsPERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy. Matematika, 1960, Nr. 3, pp. 256-260 1b

TEXT: The problem

$$\frac{d^2y(x)}{dx^2} + p_1(x) \frac{dy(x)}{dx} + p_2(x)y(x) + g_1(x) \frac{dy(x-\tau)}{dx} + g_2(x)y(x-\tau) = f(x)$$

$y(x) = \psi(x)$ for $-\tau \leq x < 0$, $y(0) = a$, $y'(0) = b$, where $p_1(x)$, $p_2(x)$, $g_1(x)$, $g_2(x)$, $f(x)$, $\psi(x)$ is continuously differentiable for $0 \leq x < +\infty$ and $\tau > 0$ is constant, is always solvable according to (Ref.1).

The approximate solution $y_N(x)$ is constructed with the aid of the functions

 $y_{k_N}(x)$, $k=0, 1, 2, \dots$ 1) $y_{0_N}(x)$ is an approximate solution of

Card 1/4

S/140/60/000/003/011/011
C111/C222

Approximate Solution of Differential Equations With a Lagging Argument With
the Aid of the Method of Oscillating Functions

$$\frac{d^2 y_0(x)}{dx^2} + p_1(x) \frac{dy_0(x)}{dx} + p_2(x) y_0(x) + g_1(x) \varphi(x-\tau) + g_2(x) \varphi(x-\tau) = f(x)$$

on $[0, \tau]$ under the conditions $y_0(0) = a$, $y'_0(0) = b$.

2) $y_{k_N}(x)$ ($k=1, 2, 3, \dots$) are approximate solutions of

$$\frac{d^2 y_k(x)}{dx^2} + p_1(x) \frac{dy_k(x)}{dx} + p_2(x) y_k(x) + g_1(x) \frac{dy_{(k-1)_N}(x-\tau)}{dx} +$$

$$+ g_2(x) y_{(k-1)_N}(x-\tau) = f(x)$$

on the $[k\tau, (k+1)\tau]$ ($k=1, 2, 3, \dots$) under the conditions $y_k(k\tau) = y_{(k-1)_N}(k\tau)$,

$y'_k(k\tau) = y'_{(k-1)_N}(k\tau)$, where $y_{(k-1)_N}(x)$ is assumed to be found. The
approximate solutions $y_{k_N}(x)$ are determined with the method of oscillating

Card 2/4

S/140/60/000/003/011/011
C111/C222

Approximate Solution of Differential Equations With a Lagging Argument With
the Aid of the Method of Oscillating Functions

functions (Ref.2).

3) Now it is put

$$y_N(x) = y_{k_N}(x) \quad (k=0, 1, 2, \dots) \text{ for } k\tau < x < (k+1)\tau.$$

The error estimation is given in agreement with (Ref.2). A numbrical example is calculated.

The same method is applied to the problem:

$$y''(x) + p_1(x)y'(x) + p_2(x)y(x) + g_1(x)y'(x-\tau) + g_2(x)y(x-\tau) = f(x)$$

$$y(x) = \varphi(x) \text{ for } -\tau \leq x < 0$$

$$u_1(y) = \alpha_1 y(0) + \alpha_1' y'(0) + \beta_1 y(1) + \beta_1' y'(1) = 0$$

$$u_2(y) = \alpha_2 y(0) + \alpha_2' y'(0) + \beta_2 y(1) + \beta_2' y'(1) = 0$$

where $p_1(x)$, $p_2(x)$, $g_1(x)$, $g_2(x)$, $f(x)$ are continuously differentiable functions on $[0, 1]$, τ is a constant positive lag, the rank of the matrix

Card 3/4

S/140/60/000/003/011/011
C111/C222

Approximate Solution of Differential Equations With a Lagging Argument With
the Aid of the Method of Oscillating Functions

$\begin{pmatrix} \alpha_1 & \alpha_1 & \beta_1 & \beta_1 \\ \alpha_2 & \alpha_2 & \beta_2 & \beta_2 \end{pmatrix}$ equals 2 and $\lambda=0$ is no eigenvalue. ✓

There are 3 Soviet references.

[Abstracter's note: The method of oscillating functions of (Ref.2) is not
described in the paper. (Ref.2) is a paper of S.I.Mel'nik in Uch.zap.
Molotovsk. un-ta, 1955, Vol.9, No.4, pp.15-24]

ASSOCIATION: Permskiy gosudarstvennyy universitet imeni A.M.Gor'kogo
(Perm State University imeni A.M.Gor'kiy)

SUBMITTED: September 16, 1958

Card 4/4

26193

S/140/60/000/005/020/021
C111/C22216.3410 U
AUTHOR: Sharkova, N.V.TITLE: Reduction of a Boundary Value Problem to a Problem With Cauchy
Initial ConditionsPERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy. Matematika, 1960,
No. 5, pp. 195 - 198

TEXT: The author considers

$$(1) \quad y'' = p(x)y' + g(x)y + r(x)$$

with $y(0) = y(1) = 0$. Let $p(x)$, $g(x)$, $r(x)$ be continuously differentiable
on $[0, 1]$; Let $\lambda = 0$ be no eigen value.

Let $y_{1N}(x)$ and $y_{2N}(x)$ be approximate solutions of (1) with the con-

ditions $y(0) = 0$, $y'(0) = b_1$ and $y(0) = 0$, $y'(0) = b_2$, where b_1 and

b_2 are chosen so that

Card 1/3

Reduction of a Boundary Value Problem to a
Problem With Cauchy Initial Conditions

86103
S/140/60/000/005/020/021
C111/C222

$$y_{1_N}(1) = \epsilon_1 > 0 \quad \epsilon_1 > M_1 h^2 \geq |y_1(x) - y_{1_N}(x)|$$

$$y_{2_N}(1) = \epsilon_2 < 0 \quad \epsilon_2 > M_2 h^2 \geq |y_2(x) - y_{2_N}(x)|$$

✓

Here $M_1 h^2$ and $M_2 h^2$ are the estimations of the error arising at the approximative solution with the aid of the method of oscillating functions. The M_1 and M_2 can be expressed by the norms of the reversion operators of

Volterra equations (cf. (Ref. 1)),
Theorem 1 : Let the curves $y_1(x)$ and $y_2(x)$ be solutions of (1) with the conditions $y(0) = 0$, $y'(0) = b_1$ and $y(0) = 0$, $y'(0) = b_2$. If they do not intersect $y_1(x)$ and $y_2(x)$ on $[0,1]$ then $y(x)$ lies between them.

Theorem 2 : If $\lambda = 0$ is no eigenvalue of the given boundary value problem but an eigenvalue of (1) for the conditions $y(0) = 0$, $y(x_i) = \alpha_i$ for $x \in [0, x_i]$, where $x_i < 1$ ($i = 0, 1, 2, \dots, n$) then the curves $y_1(x)$ and

Card 2/3

06195

Reduction of a Boundary Value Problem to a
Problem With Cauchy Initial Conditions

S/140/60/000/005/020/021
C111/C222

$y_2(x)$ intersect in all points x_i and only in them.

Theorem 3 : If $\lambda = 0$ is no eigenvalue of the given boundary value problem
then $y(x)$ lies between $y_1(x)$ and $y_2(x)$ for $x \in [0, 1]$.

According to the theorems, the author recommends one of the curves $y_1(x)$
or $y_2(x)$ as an approximate solution. The error is e.g. for $y_1(x)$:

$$|y(x) - y_1(x)| < \max_{0 \leq x \leq 1} |y_1(x) - y_2(x)| + \max(M_1, M_2) \cdot h^2$$

There is 1 Soviet reference.

[Abstracter's note : (Ref. 1) is a paper of S.I. Mel'nik in Uch. zap.
Permsk un - ta, 1955, Vol. 9, No. 4, pp. 15 - 24]

ASSOCIATION: Permskiy gosudarstvennyy universitet imeni A.M. Gor'kogo
(Perm' State University imeni A.M. Gor'kiy)

SUBMITTED: September 26, 1958

Card 3/3

L 45185-66 EWT(d)/T IJP(c)
ACC NR: AR6027459 SOURCE CODE: UR/0044/66/000/005/B044/B044

AUTHOR: Sharkova, N. V.

28
B

ORG: none

TITLE: Approximate solution of differential equations with a deflected argument
by a method of oscillating functions /6

SOURCE: Ref. zh. Matematika, Abs. 5B209

REF SOURCE: Uch. zap. Permsk. un-t, no. 103, 1963, 85-102

TOPIC TAGS: approximate solution, differential equation, function analysis,
oscillating function

ABSTRACT: A method of oscillating functions, proposed by S. I. Mel'nik
(RZhMat, 1954, 5256; 1957, 5942), is applied to the solution of differential
equations with a delayed argument, neutral-type equations, and equations with a
leading argument. The results obtained previously by the author (RZhMat, 1961,
10B220), are improved in this study. Summary. [Translation of abstract] [NT]

SUB CODE: 12/
Card 1/1 *plw*

UDC: 517.949.2

S/140/63/000/001/004/006
E032/E314

AUTHOR: Sharkova, N.V.

TITLE: Approximate solution of integral-differential equations and differential equations with retarded argument by the method of oscillating functions

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy, Matematika, no. 1, 1963, 158 - 168

TEXT: The method of oscillating functions put forward by S.I. Mel'nik (DAN SSSR, V.XCV, no. 4, 1954; Uchen. zap. Permsk. Gosn-ta, v.9, no. 4, 1955) is applied to the approximate solution of integral differential equations and linear and nonlinear differential equations with retarded argument (variable retardation). The integral differential equation considered is

$$y'(x) = \int_0^{\infty} y(x-s) dr(x, s) + f(x) \quad (1)$$

where $f(x)$ is continuous for $0 \leq x \leq \ell$ and the kernel $r(x, s)$ is defined for $0 \leq x \leq \ell$, $0 \leq s \leq \infty$. Next, it is
Card 1/6

S/140/63/000/001/004/006
E032/E514

Approximate solution

assumed that for $0 \leq s \leq l$ there exist continuous functions $\sigma(s \leq 0)$, $V(x \geq 0)$ and $M(x)$, which are such that $r(x, 0) \equiv 0$,
 $\sigma(x)$
 $\int_0^{\infty} r(x, s) \leq V(s)$, $\int_0^{\infty} r(x, s) = M(x)r(x, s) \equiv r(x, \sigma(x))$ when
 $s=0$
 $\sigma(x) < s < +\infty$ and $r(x, s)$ is such that for any $x \in [0, l]$

$$\lim_{\substack{t \rightarrow x \\ 0 \leq t \leq l}} \int_0^{\sigma(x)+1} |r(t, s) - r(x, s)| ds = 0 .$$

It is further supposed that $y(0) = \varphi(0)$, $y(x - s) = \varphi(x - s)$,
when $x - s < 0$, where $\varphi(x)$ is continuous for $-\infty < x \leq 0$
and $|\varphi(x)| \leq K$, $|\varphi(0)| < K$. The approximate solution of Eq.(1)
is sought in the form

Card 2/6

S/140/63/000/001/004/006
E032/E314

Approximate solution

$$y_{N_k}(x) = a_k(x - x_k) + b_k, \\ x_k \leq x < x_{k+1} \quad (k = 0, 1, 2, \dots, N-1), \quad \max |x_{k+1} - x_k| = h, \\ 0 = x_0 < x_1 < \dots < x_N = \ell.$$

The coefficient a_k are then determined by the oscillating-function method and the accuracy of the approximation is estimated. Next, an equation of the form

$$y''(x) = p_1(x)y'(x) + p_2(x)y(x) + g_1(x)y'(x - \Delta(x)) + \\ + g_2(x)y(x - \Delta(x)) + f(x) \quad (9)$$

is considered subject to the conditions

$$y(0) = \varphi(0) = a, \quad y'(0) = b, \quad y(x - \Delta(x)) \equiv \varphi(x - \Delta(x))$$

Card 3/6

S/140/65/000/001/004/006
E032/E314

Approximate solution

when $x - \Delta(x) < 0$, where $p_1(x)$, $p_2(x)$, $g_1(x)$, $g_2(x)$ and $f(x)$ are differentiable in $0 \leq x \leq \ell$, $\varphi(x)$ is continuously differentiable in $-\infty < x \leq 0$ and $\Delta(x) \geq 0$ is continuous in $0 \leq x \leq \ell$. The approximate solution of this problem is sought in the form

$$y_{N_k}(x) = \frac{a_k}{2} (x - x_k)^2 + b_k(x - x_k) + c_k,$$

$$x_k < x < x_{k+1} \quad (k = 0, 1, 2, \dots, N-1), \quad \max |x_{k+1} - x_k| = h$$

$$0 = x_0 < x_1 < \dots < x_N = \ell.$$

Again, the coefficients are evaluated by the oscillating-function method and the errors involved are estimated. Finally, the same methods are used to solve the equation

$$y'' = f(x, y)(x), \quad y'(x), \quad y(x - \Delta(x)), \quad y'(x - \Delta(x))) \quad (15)$$

Card 4/6

S/140/65/000/001/004/006
E052/E314

Approximate solution

subject to the conditions $y(0) = \varphi(0) = a$, $y'(0) = b$,
 $y(x - \Delta(x)) \leq \varphi(x - \Delta(x))$, when $x - \Delta(x) > 0$, where f is
continuous with respect to all its arguments in $0 \leq x \leq \ell$,
 $-\infty < y$, $y' < +\infty$ and satisfy the Lipschitz condition for
all its arguments beginning with the second, $\varphi(x)$ is an
initial function which is continuously differentiable in $-\infty < x \leq$
 ≤ 0 and $\Delta(x) \geq 0$ is continuous in $0 \leq x \leq \ell$. Eq. (15)
is equivalent to (subject to the given initial conditions) the
following system of second-order equations

$$\begin{aligned} y'(x) &= z(x), \\ z'(x) &= f(x, y(x), z(x), y(x - \Delta(x)), z(x - \Delta(x))) \end{aligned} \quad (16)$$

subject to the conditions $y(0) = a$, $z(0) = b$, $y(x - \Delta(x)) \leq$
 $\leq \varphi(x - \Delta(x))$ when $x - \Delta(x) < 0$. The approximate solution
of Eq. (16) is then seen to be in the form

$$y_{N_k}(x) = a_k(x - x_k) + b_k, \quad z_{N_k}(x) = A_k(x - x_k) + B_k, \text{ where}$$

Card 5/6

Approximate aolution

S/140/63/000/001/004/006
E032/E314

$x_k \leq x \leq x_{k+1}$ ($k = 0, 1, 2, \dots, N - 1$), $\max |x_{k+1} - x_k| = h$,
 $0 = x_0 < x_1 < \dots < x_N = \ell$

and the accuracy of the approximation is estimated.

ASSOCIATION: Permskiy gosudarstvennyy universitet im.
A.M. Gor'kogo (Perm. State University im.
A.M. Gor'kiy)

SUBMITTED: December 30, 1959

Card 6/6

L 11113-63

EWT(d)/FCC(w)/BDS--AFFTC--Pg-4--IJP(C)

ACCESSION NR: AP3000958

S/0140/63/000/003/0173/0184

AUTHOR: Sharkova, N. V. (Perm')

55

TITLE: The approximate solution of differential equations with a deviating argument

SOURCE: IVUZ. Matematika, no. 3, 1963, 173-184

TOPIC TAGS: differential equation, deviating argument, approximate solution, oscillatory function method

ABSTRACT: The method of oscillatory functions proposed by S. I. Mel'nik for the solution of differential and integral equations is applied to the approximate solution of second-order differential equations with a deviating argument. The following four cases are investigated: a) linear equations of the neutral type; b) nonlinear equations of the neutral type; c) linear equations with a leading argument; and d) nonlinear equations with a leading argument. The approximate solutions of types (a), (c), and (d) are sought in the form

$$Y_{N_k}(x) = \frac{a_k}{2}(x - x_k)^2 + b_k(x - x_k) + c_k, \quad (1)$$

Card 1/2

L 11113-63
ACCESSION NR: AP3000958

where $x_k \leq x \leq x_{k+1}$ and $\max(x_{k+1} - x_k) = h$. Constants $c_k, b_k, (k = 1, 2, \dots, N - 1)$ are determined from the conditions of continuity of the approximate solution and of its first derivative at the junction points of two parabolas; constants a_k are determined from the oscillatory function. In case (b) a system equivalent to two first-order differential equations is derived, and the approximate solution is sought in the form

$$\begin{aligned} Y_{N_k}(x) &= a_k(x - x_k) + b_k \\ Z_{N_k}(x) &= A_k(x - x_k) + B_k \end{aligned} \quad (2)$$

The method of determining the constants a_k, b_k, A_k , and B_k is presented. An estimate of the error of the approximation is derived: In case (a) it is proportional to h^3 ; in case (b), to h^2 ; and in the last two cases, to h . Orig. art. has: 19 equations.

ASSOCIATION: none

SUBMITTED: 03Jun60

DATE ACQ: 12Jun63

ENCL: 00

SUB CODE: MM

NO REF Sov: 004

OTHER: 000

Card 2/2 ch *www*

SHARKOVA, T.T.

Some new species of the Silurian and Devonian Tabulata of
southeastern Kazakhstan. Paleont. zhur. no. 1:20-25 '64.
(MIRA 17:7)

1. Vsesoyuznyy aerogeologicheskiy trest.

SHARKOVA, T.T.

New genus Axuolites (Favostida) from the Upper Silurian in
Kazakhstan. Paleot. zhur. no.3:117-119 '63. (MIRA 16:10)

1. Vsesoyuznyy aerogeologicheskiy trest.

KRUT', I.V.; YAKOVLEV, L.I.; KROPACHEV, S.M.; LYASHENKO, A.I.;
SHARKOVA, T.T.

Stratigraphic position and structure of the Karashay series
in the Northern Caucasus. Izv. AN SSSR. Ser. geol. 28 no.10:
49-59 0 '63. (MIRA 16:11)

1. Tsentral'nyy nauchno-issledovatel'skiy geologorazvedochnyy
institut, Moskva.

SHARKOVA, V.M.

Use of the shaft-type electric furnace instead of the chamber-type
furnace for heat treating diesel piston rings. Prom. energ. 11 no.10:
26 0 '56. (MIRA 9:11)

(Electric furnaces) (Piston rings)

SHARKOVA, V. V.

N/5
621.212
.A3

ADADEMIYA NAUK SSSR. LABORATORIYA AEROMETODOV. PRIMENENIYE AEROMETODOV
Dlya GEOFIZICHESKIH ISSEKOVANIY MORSKOGO DNA, KRATKOYE METODICHESKYE PUKO-
VODSTVO (USE OF AERO-METHODS FOR UNDERWATER GEOLOGICAL SURVEYS) POD RED. V. V.
SHARKOVA I D. M. KUDRITSKOGO. LENINGRAD, GOSTCPTEKHIZDAT, 1956. 254, (2) p.
ILLUS., DIAGRS., TABLES. "LITERATURA": p. 252-(255)

SHARKOVA, V.Ye.

Calcium arsenite is toxic to the shield bug Eurygaster integriceps.
Zashch.rast.ct vred.i bol. 4 no.6:37-38 N-D '59. (MIRA 15:11)

1. Nachal'nik proizvodstvennogo uchastka Rostovskogo otryada po
bor'be s vreditelyami i boleznyami rasteniy.
(Rostov Province--Eurygasters--Extermination) (Calcium arsenites)

SHARKOVA, V.Ye.

They reduced the costs. Zashch.rast.ot vred.i bol. 5 no.3:6-7
Mr '60. (MIRA 16:1)

1. Nachal'nik proizvodstvennogo uchastka Rostovskogo otryada
po bor'be s vreditelyami i boleznyami rasteniy.
(Rostov Province—Eurygasters—Extermination)

ARKHANGEL'SKIY, N.I.; SHTEYNBERG, G.G., nauchnyy sotrudnik; SHARKOVA, V.Ye.

Poisoned objects providing cover as a method for controlling
injurious insects. Zashch. rast. ot vred. i bol. 5 no.9:27-32
(MIRA 15:6)
S '60.

1. Chlen-korrespondent Vsesoyuznnyy akademii sel'skokhozyaystvennykh
nauk imeni V.I. Lenina (for Arkhangel'skiy). 2. Nachal'nik
uchastka Rostovskogo otchela bor'by s vreditelyami sel'skogo
khozyaystva (for Sharkova).
(Insecticides)

SHARKOVA, V. Ye.

Socialist obligations are being fulfilled. Zashch. rast. ot
vred. i bol. 5 no.10:4-6 0 '60. (MIRA 16:1)

1. Nachal'nik proizvodstvennogo uchastka Rostovskogo oblastnogo
otryada po bor'be s vreditelyami i boleznyami rasteniy.

(Roston Province--Plants, Protection of)

SHARKOVA, V.Ye.

One of the best. Zashch. rast. ot vred. i bol. 6 no.8:42-43
Ag '61. (MIRA 15:12)

1. Obshchestvennyy korrespondent zhurnala "Zashchita rasteniy".
(Plants, Protection of)

SHARKOVA, V.Ye.

Each experimental demonstration farm should be provided with
an agronomist for plant protection. Zashch. rast. ot vred. i
bol. 6 no.11:7-8 N '61. (MIRA 16:4)

1. Nachal'nik otdela zashchity rasteniy Rostovskogo oblastnogo
upravleniya sel'skogo khozyaystva.
(Rostov Province—Plants, Protection of)

YAREMENKO, N.Ye.; SHARKOVA, V.Ye.

Protecting grain with arsenic preparations. Zashch. rast. ot
vred. i bol. 7 no.12:21-23 D '62. (MIRA 16:7)

1. Glavnyy agronom Rostovskogo otryada po bor'be s vreditelyami
i boleznyami rasteniy (for Yaremenko). 2. Starshiy agronom
Rostovskogo otryada po bor'be s vreditelyami i boleznyami
rasteniy (for Sharkova).
(Rostov Province---Rodent control) (Arsenic)

SHARKOVA, V.Ye.

On the "TSelinskii" State Farm. Zashch. rast. ot vred. i bol.
9 no.8;9-10 '64. (MIRA 17;12)

1. Starshiy agronom Rostovskoy stantsii zashchity rasteniy.

SHAROV, A. I., VIREL, A. A., BOGOVIN, A. A.; Printmaja uchastvuyet
SHVYAKOV, K. P.

Polymer chain conversions of a graft copolymer of cellulose
and *p*-glycidyl methacrylate. Vysokomol. soed. 6 no. 5:951-955
(MIRA 17:6)

1. Post-apply coating, inventors.

L 1354-66 EPF(c)/EWT(m)/ETC/EWG(m)/T/EWP(j) RPL DS/WW/RM
ACCESSION NR: AP5024397 UR/0286/65/000/015/0080/0080
678.542
678.744.322-13 32
B

AUTHOR: Rogovin, Z. A.; Vurnik, A. D.; Sharkova, Ye. F. 44.55 15

TITLE: A method for producing a graft copolymer. 9.44.55 Class 39, No. 173404

SOURCE: Byulleten' izobreteniy i tovarnykh znakov, no. 15, 1965, 80

TOPIC TAGS: graft copolymer, cellulose plastic, acrylic plastic, methacrylate plastic, ion exchange resin, radical polymerization

ABSTRACT: This Author's Certificate introduces: 1. A method for producing a graft copolymer by radical graft copolymerization of cellulose and an ester of acrylic or methacrylic acid. A wider selection of monomers which can be grafted to cellulose is provided by using glycidylacrylate or glycidylmethacrylate. 2. A modification of this method in which an ion-exchange copolymer is produced by treating the finished graft copolymer in compounds which react with its α -oxide cycles, e. g. aqueous solutions of primary or secondary amines, sulfite or bisulfite of sodium.

ASSOCIATION: none

SUBMITTED: 17Jun63

NO REF SOV: 000

ENCL: 00 SUB CODE: MT, GC

OTHER: 000

K1
Card 1/1

BALABAYEVA, M.D.; SHARKOVA, Ye.F.; ZHBANKOV, R.G.; VIRNIK, A.D.; ROGOVIN, Z.A.

Infrared spectroscopy method of studying the structure of
some graft copolymers of cellulose. Vysokom. soed. 7
no.10:1763-1766 0 '65. (MIRA 18:11)

1. Moskovskiy tekstil'nyy institut.

SHARKWA, H. S., J. R. LEE, A. G. COLE, and J. R. COLE

Министерство труда и социальной политики Российской Федерации

APPROVED FOR RELEASE: 08/23/2000

CIA-RDP86-00513R001548620004-1"

SEVERIN, S.Ye.; TELEPNEVA, V.I.; ZAYTSEVA, N.N.; SHARKOVA, Ye.V.

Particular features of oxidative phosphorylation in denervated
skeletal muscles. Ukr.biokhim.zhur. 37 no.5:787-797 '65.
(MIRA 18:10)

1. Kafedra biokhimii zhivotnykh Moskovskogo gosudarstvennogo
universiteta.

AYNBINDER, N.M.; DIL'MAN, V.M.; MUKHINA, Ye.P.; NECHAYEVA, I.D.; SHARKOVA,
Zh.M.

Experience with the antibiotic 2703 in six patients with chorio-
epithelioma of the uterus. Vop. onk. 10 no.5:103-107 '64.
(MIRA 18:8)

1. Iz Instituta onkologii AMN SSSR (dir. - prof. A.I.Serebrov).
Adres avtorov: Leningrad, P-129, 2-ya Berezovaya alleya, 3,
Institut onkologii AMN SSSR.